

# SCHIZOPHRENIA

## Let's talk about it

Schizophrenia is a serious mental illness and one of the top 15 leading causes of disability worldwide. The disorder is characterized by alterations to a person's thoughts, feelings, and behaviors, which can include a loss of contact with reality known as psychosis. These symptoms typically emerge in adolescence or early adulthood and can be persistent and disabling when left untreated, interfering with a person's ability to engage in typical school, work, and social activities.

### What is it?

Schizophrenia is a serious mental illness that affects how a person thinks, feels, and behaves. People with schizophrenia may seem as though they have lost touch with reality, which can be distressing for them and for their family and friends. The symptoms of schizophrenia can make it difficult to participate in usual, everyday activities, but effective treatments are available. Many people who receive treatment can engage in school or work, achieve independence, and enjoy personal relationships.

### What causes it?

Several factors may contribute to a person's risk of developing schizophrenia, including:

- **Genetics.** Schizophrenia sometimes runs in families. However, just because one family member has schizophrenia, it does not mean that other members of the family also will have it. Studies suggest that many different genes may increase a person's chances of developing schizophrenia, but that no single gene causes the disorder by itself.
- **Environment.** Research suggests that a combination of genetic factors and aspects of a person's environment and life experiences may play a role in the development of schizophrenia. These environmental factors may include living in poverty, stressful or dangerous surroundings, and exposure to viruses or nutritional problems before birth.
- **Brain structure and function.** Research shows that people with schizophrenia may be more likely to have differences in the size of certain brain areas and in connections between brain areas. Some of these brain differences may develop before birth. Researchers are working to better understand how brain structure and function may relate to schizophrenia.

### Signs & Symptoms

It's important to recognize the symptoms of schizophrenia and seek help as early as possible. People with schizophrenia are usually diagnosed between the ages of 16 and 30, after the first episode of psychosis. Starting treatment as soon as possible following the first episode of psychosis is an important step toward recovery. However, research shows that gradual changes in thinking, mood, and social functioning often appear before the first episode of psychosis. Schizophrenia is rare in younger children.

Schizophrenia symptoms can differ from person to person, but they generally fall into three main categories: psychotic, negative, and cognitive.

#### Psychotic symptoms

Psychotic symptoms include changes in the way a person thinks, acts, and experiences the world. People with psychotic symptoms may lose a shared sense of reality with others and experience the world in a distorted way. For some people, these symptoms come and go. For others, the symptoms become stable over time. Psychotic symptoms include:

- **Hallucinations:** When a person sees, hears, smells, tastes, or feels things that are not actually there. Hearing voices is common for people with schizophrenia. People who hear voices may hear them for a long time before family or friends notice a problem.
- **Delusions:** When a person has strong beliefs that are not true and may seem irrational to others. For example, individuals experiencing delusions may believe that people on the radio and television are sending special messages that require a certain response, or they may believe that they are in danger or that others are trying to hurt them

## FOR MORE INFORMATION CONTACT US

Iroquois Center for Human Development  
(620) 723 - 2272

Source: National Institute of Mental Health



# SCHIZOPHRENIA

## Signs & Symptoms (cont.)

- **Thought disorder:** When a person has ways of thinking that are unusual or illogical. People with thought disorder may have trouble organizing their thoughts and speech. Sometimes a person will stop talking in the middle of a thought, jump from topic to topic, or make up words that have no meaning.
- **Movement disorder:** When a person exhibits abnormal body movements. People with movement disorder may repeat certain motions over and over.

### Negative symptoms

Negative symptoms include loss of motivation, loss of interest or enjoyment in daily activities, withdrawal from social life, difficulty showing emotions, and difficulty functioning normally. Negative symptoms include:

- Having trouble planning and sticking with activities, such as grocery shopping
- Having trouble anticipating and feeling pleasure in everyday life
- Talking in a dull voice and showing limited facial expression
- Avoiding social interaction or interacting in socially awkward ways
- Having very low energy and spending a lot of time in passive activities. In extreme cases, a person might stop moving or talking for a while, which is a rare condition called catatonia.

These symptoms are sometimes mistaken for symptoms of depression or other mental illnesses.

### Cognitive symptoms

Cognitive symptoms include problems with attention, concentration, and memory. These symptoms can make it hard to follow a conversation, learn new things, or remember appointments. A person's level of cognitive functioning is one of the best predictors of their day-to-day functioning. Cognitive functioning is evaluated using specific tests. Cognitive symptoms include:

- Having trouble processing information to make decisions
- Having trouble using information immediately after learning it
- Having trouble focusing or paying attention

### Risk of violence

Most people with schizophrenia are not violent. Overall, people with schizophrenia are more likely than those without the illness to be harmed by others. For people with schizophrenia, the risk of self-harm and of violence to others is greatest when the illness is untreated. It is important to help people who are showing symptoms to get treatment as quickly as possible.

### Schizophrenia vs. dissociative identity disorder

Although some of the signs may seem similar on the surface, schizophrenia is not dissociative identity disorder (which used to be called multiple personality disorder or split personality). People with dissociative identity disorder have two or more distinct identities that are present and that alternately take control of them.

## FOR MORE INFORMATION CONTACT US

Iroquois Center for Human Development  
(620) 723 - 2272

Source: National Institute of Mental Health



# SCHIZOPHRENIA

## How is it treated?

Current treatments for schizophrenia focus on helping individuals manage their symptoms, improve day-to-day functioning, and achieve personal life goals, such as completing education, pursuing a career, and having fulfilling relationships.

### Antipsychotic medications

Antipsychotic medications can help make psychotic symptoms less intense and less frequent. These medications are usually taken every day in a pill or liquid form. Some antipsychotic medications are given as injections once or twice a month. If a person's symptoms do not improve with usual antipsychotic medications, they may be prescribed clozapine. People who take clozapine must have regular blood tests to check for a potentially dangerous side effect that occurs in 1% to 2% of patients. People respond to antipsychotic medications in different ways. It is important to report any side effects to a health care provider. Many people taking antipsychotic medications experience side effects such as weight gain, dry mouth, restlessness, and drowsiness when they start taking these medications. Some of these side effects may go away over time, while others may last.

### Psychosocial treatments

Psychosocial treatments help people find solutions to everyday challenges and manage symptoms while attending school, working, and forming relationships. These treatments are often used together with antipsychotic medication. People who participate in regular psychosocial treatment are less likely to have symptoms reoccur or to be hospitalized. Examples of this kind of treatment include cognitive behavioral therapy, behavioral skills training, supported employment, and cognitive remediation interventions.

### Coordinated specialty care

Coordinated specialty care (CSC) programs are recovery-focused programs for people with first episode psychosis, an early stage of schizophrenia. Health professionals and specialists work together as a team to provide CSC, which includes psychotherapy, medication, case management, employment and education support, and family education and support. The treatment team works collaboratively with the individual to make treatment decisions, involving family members as much as possible.

Compared with typical care, CSC is more effective in reducing symptoms, improving quality of life, and increasing involvement in work or school.

## FOR MORE INFORMATION CONTACT US

Iroquois Center for Human Development  
(620) 723 - 2272

Source: National Institute of Mental Health